

Comment on “Reaction Coordinates and Pathways of Mechanochemical Transformations”

Wolfgang Quapp^{*,†} and Josep Maria Bofill^{*,‡}

*Mathematisches Institut, Universität Leipzig, PF 100920, D-04009 Leipzig, Germany, and
Departament de Química Orgànica, Universitat de Barcelona, and Institut de Química
Teòrica i Computacional, Universitat de Barcelona, (IQTCUB), Martí i Franquès, 1,
08028 Barcelona Spain*

E-mail: quapp@uni-leipzig.de; jmb@ub.edu

*To whom correspondence should be addressed

†Leipzig University

‡Universitat de Barcelona

The adiabatic potential energy surface (PES) is a basic concept of many theoretical chemistry models. In the last years, the phenomena of the action of a mechanical stress over a molecular system have motivated experimental and theoretical researches. In a recent article, Avdoshenko and Makarov¹ describe how the concepts of an effective PES and of a reaction path (RP), or a reaction coordinate can be used for mechanochemistry.

The RP is a one-dimensional description of a pathway on the PES in an N -dimensional configuration space. We use $N = 3n - 6$ non-redundant internal coordinates, and n is the number of the atoms of the molecule. An early used curve is the distinguished coordinate,² which was later generalized as a distinguished coordinate path (DCP)³ and finally refined as Newton trajectory (NT).⁴⁻⁶ For this type of RP holds the following property: the gradient of the PES points into the same direction at every point of the curve. It is the reason that NTs should be taken into account for mechanochemical problems.

Basically the mechanochemistry model¹ consists in a first order perturbation on the associated PES of the unperturbed molecular system due to a stress or pulling force, \mathbf{f}

$$V_f(\mathbf{r}) = V(\mathbf{r}) - \mathbf{f}^T \mathbf{R} . \quad (1)$$

\mathbf{R} is the distance between the two pulling points of the molecule.⁷ It will be associated with one of the coordinates,⁷ or a linear combination of them. So we can assume that $dR \approx dr$ for a coordinate change in direction of \mathbf{R} . The potential $V_f(\mathbf{r})$ with a fixed \mathbf{f} can be seen as an effective PES where 'normal' chemistry takes place. Due to the external force, the stationary points of $V_f(\mathbf{r})$ are located at different positions,⁸ with respect to the unperturbed potential. For the new minimum holds with Eq.(1)

$$\nabla_{\mathbf{r}} V_f(\mathbf{r}) = \mathbf{0} = \mathbf{g} - \mathbf{f} , \quad (2)$$

thus one searches a point where the gradient, \mathbf{g} , of the zero-force PES has to be equal to the mechanochemical force. If the mechanical stress in a defined direction is $\mathbf{f} = F\mathbf{l}$ with a

fixed unit vector, \mathbf{l} , then it is $\mathbf{l} = \mathbf{g}/|\mathbf{g}|$ and $F = |\mathbf{g}|$. Another form of Eq.(2) is the projector equation which was applied many years ago^{5,9}

$$(\mathbf{U} - \mathbf{l}\mathbf{l}^T) \mathbf{g} = \mathbf{0} \quad (3)$$

where \mathbf{U} is the unit matrix. Solution curves of both Eqs. (2) and (3) are equal, and they are also equal⁵ to the solutions of the differential equation of Branin¹⁰

$$\frac{d\mathbf{r}}{dt} = \pm \mathbf{A}(\mathbf{r}) \mathbf{g}(\mathbf{r}) . \quad (4)$$

t is a curve parameter and the matrix \mathbf{A} is constructed by a multiplication of the inverse Hessian with the determinant of the Hessian, $\mathbf{A} = \text{Det}(\mathbf{H}) \mathbf{H}^{-1}$. Curves $\mathbf{r}(t)$ satisfying (4) are called Newton trajectories (NT). The name is also used for the equivalent solution curves of the Eqs. (2) and (3).

For different, point-to-point changing forces, F , one should get a curve of the 'reaction path following force displaced stationary points (FDSPs)^{1,7,11}

$$\delta\mathbf{r} = \mathbf{H}^{-1}(\mathbf{r}) \mathbf{f}(\mathbf{r}) = \mathbf{H}^{-1}(\mathbf{r}) \mathbf{g}(\mathbf{r}) \quad (5)$$

where $\delta\mathbf{r}$ is the distance from the minimum of $V(\mathbf{r})$ to the minimum of $V_f(\mathbf{r})$. Note that the inverse Hessian \mathbf{H}^{-1} will be singular on the pathway from the minimum to the saddle point of the original PES, and Eq.(5) loses its meaning. However, such points are the important catastrophe points in ref. 1.

There is another way to make the FDSPs without the singularity: the Branin Eq. (4). It is the result of a different parametrization of the curve parameter, t , by the multiplication of the right hand side of Eq.(5) by the determinant of the Hessian. The determinant is a number. It does not change the direction of the vector of the right hand side of Eq. (5).

However, it removes the singularity of \mathbf{H}^{-1} on the way to the SP. The matrix \mathbf{A} is named the desingularized inverse Hessian, or the adjoint to \mathbf{H} .

The solution curve of the Branin equation to a given initial direction is a regular curve (if no valley-ridge inflection point is crossed^{5,12} – but this is another, seldom property). The Branin equation is a well-know model for RPs.⁵ However, the model is used here for the FDSPs. The stationary points of the different effective potentials with fixed \mathbf{l} move with increasing F on the original PES along an NT. The behaviour of NTs is well known.¹³

We will still discuss an example:¹ the ring opening of *trans*-1,2-dimethylcyclobutane. There is chosen a DCP for the curve of FDSPs. However, the DCP jumps over the PES. The DCP method has been criticized during its first years for such jumps.³ Avdoshenko and Makarov¹ circumvent the problem by calculating two different curves, one DCP from the minimum, and one direct FDSPs by Eq. (5) from the SP, and both curves meet at the jump point. The reason is, both curves are parts of a Newton trajectory (NT) with a turning point, compare Fig. 1 and Fig. 2 of ref. 1.




Figure 1: NTs on a PES being similar to Fig.2 in ref. 1. The bold faced NTs are direct curves from the lower minimum to the SP. The thin dashed curves are NTs with a turning point (TP) in the lower bowl with higher energy than the SP. They may be questionable pulling scenarios. The thick dashed NT goes wrong. It represents a pulling force which does not enforce the desired reaction.

A general solution of the DCP jump problems was given in 1998 by Quapp et al.^{4,5} using the Eqs. (3) and (4). NTs can have turning points (TPs) where the energy profile over the curve changes its direction. These TPs are the break-down points of the DCP method. Important points are also the valley-ridge inflection points (VRI) of the PES^{5,12} where singular NTs bifurcate. VRIs discriminate different families of NTs which lead to different SPs around the initial minimum. An NT which leads to a not desired SP represents an uncorrect pulling scenario. On the PES of Fig. 1 a VRI is at $\approx(3,0)$. We represent an NT (thick and dashed) which does not find the SP.

It should be noted that a further mathematical method gives curves in the coordinate space which are equivalent to NTs, the Newton homotopy method.¹⁴⁻¹⁶ It can extend the arsenal of methods to get the FDSPs curve.

We conclude that Newton trajectories can be used for the 'reaction path following force displaced stationary points' (FDSPs). This kind of curves form an important model for the treatment of mechanochemistry. The theory of NTs is well prepared. It is to hope that it can accomplish deeper insights into the understanding of mechanochemistry.

References

- (1) Avdoshenko, S. M.; Makarov, D. E. Reaction Coordinates and Pathways of Mechanochemical Transformations. *J. Phys. Chem. B* **2015**, DOI: [10.1021/acs.jpcc.5b07613](https://doi.org/10.1021/acs.jpcc.5b07613).
- (2) Rothman, M. J.; Lohr, Jr., L. L. Analysis of an Energy Minimization Method for Locating Transition States on Potential Energy Hypersurfaces. *Chem. Phys. Lett.* **1980**, *70*, 405-409.
- (3) Williams, I. H.; Maggiora, G. M. Use and Abuse of the Distinguished-Coordinate Method For Transition-State Structure Searching. *J. Mol. Struct. (Theochem)* **1982**, *89*, 365-378.

- (4) Quapp, W.; Hirsch, M.; Imig, O.; Heidrich, D. Searching for Saddle Points of Potential Energy Surfaces by Following a Reduced Gradient. *J. Comput. Chem.* **1998**, *19*, 1087–1100.
- (5) Quapp, W.; Hirsch, M.; Heidrich, D. Bifurcation of Reaction Pathways: the Set of Valley-Ridge Inflection Points of a Simple Three-dimensional Potential Energy Surface. *Theor. Chem. Acc.* **1998**, *100*, 285–299.
- (6) Bofill, J. M.; Anglada, J. M. Finding Transition States using Reduced Potential-Energy Surfaces. *Theor. Chem. Acc.* **2001**, *105*, 463–472.
- (7) Bailey, A.; Mosey, N. J. Prediction of Reaction Barriers and Force-induced Instabilities under Mechanochemical Conditions with an Approximate Model: A Case Study of the Ring Opening of 1,3-cyclohexadiene. *J. Chem. Phys.* **2012**, *136*, 044102.
- (8) Konda, S. S. M.; Brantley, J. M.; Bielawski, C. W.; Makarov, D. E. Chemical Reactions Modulated by Mechanical Stress: Extended Bell Theory. *J. Chem. Phys.* **2011**, *135*, 164103.
- (9) Crehuet, R.; Bofill, J. M.; Anglada, J. M. A new Look at the Reduced-Gradient-Following Path. *Theor. Chem. Acc.* **2002**, *107*, 130–139.
- (10) Branin, F. Widely Convergent Methods for Finding Multiple Solutions of Simultaneous Nonlinear Equations. *IBM J Res Develop* **1972**, *16*, 504–522.
- (11) Konda, S. S. M.; Avdoshenko, S. M.; Makarov, D. E. Exploring the Topography of the Stress-modified Energy Landscapes of Mechanosensitive Molecules. *J. Chem. Phys.* **2014**, *140*, 104114.
- (12) Quapp, W. Can We Understand the Branching of Reaction Valleys for more than two Degrees of Freedom? *J. Math. Chem.* **2015**, *54*, 137 – 148.

- (13) Quapp, W. Reduced Gradient Methods and their Relation to Reaction Paths. *J. Theoret. Comput. Chem.* **2003**, *2*, 385–417.
- (14) Ackermann, S.; Kliesch, W. Computation of Stationary Points via Homotopy Method. *Theor. Chem. Acc.* **1998**, *99*, 255–264.
- (15) Mehta, D.; Chen, T.; Morgan, J. W. R.; Wales, D. J. Exploring the Potential Energy Landscape of the Thomson Problem via Newton Homotopies. *J. Chem. Phys.* **2015**, *142*, 194113.
- (16) Bofill, J. M. Comment on "Exploring the Potential Energy Landscape of the Thomson Problem via Newton Homotopies" [J. Chem. Phys. 142, 194113 (2015)]. *J. Chem. Phys.* **2015**, *143*, 247101.

Graphical TOC Entry

